New York State Regents Examination in Physical Setting/Earth Science

2016 Technical Report

Prepared for the New York State Education Department
by Pearson

March 2017

CHAPTER 1: INTRODUCTION 1

1.1 INTRODUCTION 1
1.2 PURPOSES OF THE EXAM 1
1.3 TARGET POPULATION (STANDARD 7.2) 2

CHAPTER 2: CLASSICAL ITEM STATISTICS (STANDARD 4.10) 4

2.1 ITEM DIFFICULTY 4
2.2 ITEM DISCRIMINATION 4
2.3 DISCRIMINATION ON DIFFICULTY SCATTER PLOTS 8
2.4 OBSERVATIONS AND INTERPRETATIONS .....

1.3 TARGET POPULATION (STANDARD 7.2)

The examinee population for the Regents Examination in Physical Setting/Earth Science is composed of students who have completed a course in Earth Science.

Table 1 provides a demographic breakdown of all students who took the August 2015, January 2016, and June 2016 Regents Examination in Physical Setting/Earth Science. All analyses in this report are based on the population described in Table 1. Annual Regents Examinati inagrainis6 ( )e-151 (bd1 (in A)-15-3 (st)-143 (2)-3 0 612 792 reW* nBT0 g/TT1 12 Tf88.344 609

l1ll-111 (resu)(s)-150 0 612 792 reW* nBT0 g/TT1 12 Tf452.35 622.9 T0(91)5[(ll)14 ( )-111 (New)10 pb(i ( )2s 2 cl

This chapter provides an overview of the two most familiar item-level statistics obtained from classical item analysis: item difficulty and item discrimination. The following results pertain only to the operational Regents Examination in Physical Setting/Earth Science items.

2.1 ITEM DIFFICULTY

$W WKH PRVW JHQHUDO OHYHO DQ LWHP¶V GLIILFXOW\ LV group (e.g., grade level).

In the mean score formula above, the individual item scores (x_i) are summed and then divided by the total number of students (n). For multiple-choice (MC) items, student scores are represented by 0s and 1s (0 = wrong, 1 = right). With 0 ±1 scoring, the equation above also represents the number of students correctly answering the item divided by the total number of students. Therefore, this is also the proportion correct for the item, or the p-value. In theory, p-values can range from 0.00 to 1.00 on the proportion-correct scale.[2] For example, if an MC item has a p-value of 0.89, it means that 89 percent of the students answered the item correctly.

Table 3 Constructed-

$$P_{ni}\ X \quad 1 = \frac{\exp \quad T_n \quad D_{ij}}{1 \quad \exp \quad T_n \quad D_{ij}}.$$

The Rasch model places both performance and item difficulty (estimated in terms of log-odds or logits) on the same continuum. When the model assumptions are met, the Rasch model provides estimates of examinee performance and item difficulty that are theoretically invariant acros

A parallel analysis (Horn, 1965) can be further helpful

$$P\ X_i \quad x_i, X_j \quad x_j \mid T \quad P\ X_i \quad x_i \mid T P\ X_j \quad x_j \mid T.$$

Marais and Andrich (2008) point out that local item dependence in the Rasch model can occur in two ways that may be difficult to distinguish. The first way occurs when the assumption of unidimensionality is violated. Here, other nuisance dimensions besides a dominant

This process results in a relationship between the raw scores on the form and the overall scale scores. The scale scores corresponding to each raw score are then rounded to the nearest integer for reporting on the conversion chart (posted at the close of each administration). The only exceptions are for the minimum and maximum raw scores and the raw scores that correspond to the scaled cut scores of 55, 65, and 85.

The minimum (zero) and maximum possible raw scores are assigned scale scores of 0 and 100, respectively. In the event that there are raw scores less than the maximum with scale scores that round to 100, their scale scores are set equal to 99. A similar process is followed with the minimum score; if any raw scores other than zero have scale scores that round to zero, their scale scores are instead set equal to one.

With regard to the cuts, if two or more scale scores round to 55, 65, or 85, the lowest raw V F R U H ¶ V V F D O H V F R U H L V V H W H T X D O W R R U D Q G V higher raw scores are set to 56, 66, or 86 as appropriate. If no scale score rounds to these critical cuts, then the raw score with the largest scale score that is less than the cut is set equal to the cut. The overarching principle when two raw scores both round to either scale score cut is that the lower of the raw scores is always assigned to be equal to the cut so that students are never penalized for this ambiguity.

The New York State Regents Examination in Physical Setting/Earth Science contains both a written and a lab (or performance) component. The lab component remains constant from

Test reliability is a measure of the internal consistency of a test (Cronbach, 1951). It is a measure of the extent to which the items on a test provide consistent information about student mastery of a domain. Reliability should ultimately demonstrate that examinee score estimates maximize consistency and therefore minimize error, or theoretically speaking, that examinees who take a test multiple times would get the same score each time.

According to the *Standards for Educational and Psychological Testing* [3] $ Q X P E H U R I I D F
can have significant effects on reliability/precision, and in some cases, these factors can lead
W R P L V L Q W H U S U H W D W L R Q V R I W H V W VAERA et al., 2014, pp. 38). First, W D N H Q
test length and the variability of observed scores can both influence reliability estimates. Tests with fewer items or with a lack of heterogeneity in scores tend to produce lower reliability estimates. Second, reliability is specifically concerned with random sources of error. Accordingly, the degree of inconsistency due to random to ranincon 12 Tf57.6417441 s (icallyo)4 ( )-4

error (Harvill, 1991). Additionally, the value of the SEM takes the group variation (i.e., score standard deviation) into account. Consider that a SEM of 3 on a 10-point test would be very different from a SEM of 3 on a 100-point test.

Traditional Standard Error of Measurement Confidence Intervals

The SEM is an index of the random variability in test scores reported in actual score units, which is why it has such great utility for test score users. SEMs allow statements regarding the SUHFLVLRQ RI LQGLYLGXDO WHVW VFRUHV 6(0V KHOS SODFH around observed scores, through construction of an approximate score band. Often referred to as confidence intervals, these bands are constructed by taking the observed scores, *X*, and adding or subtracting a multiplicative factor of the SEM. As an example, students with a given true score will have observed scores that fall between ±1 SEM about two-thirds of the time.[4] For ±2 SEM confidence intervals, this increases to about 95 percent.

The Coefficient Alpha and associated SEM for the Regents Examination in Physical Setting/Earth Science are provided in Table 7.

Table 7 Reliabilities and Standard Errors of Measurement: Regents Examination in Physical Setting/Earth Science

Subject	Coefficient Alpha	SEM
Physical Setting/ Earth Science	0.93	5.13

Assuming normally distributed scores, one would expect about two-thirds of the observations to be within one standard deviation of the mean. An estimate of the standard deviation of the true scores can be computed as

$$_{T} V\sqrt{\ ^{2}_{x}V\ \ ^{2}_{x}V(1\ \ U_{xx})}\ .$$

Conditional Standard Error of Measurement

Every time an assessment is administered, the score that the student receives contains some error. If the same exam were administered an infinite number of times to the same VWXGHQW WKH PHDQ RI WKH GLVWULEXWLRQ RI WKH VWXGHQW¶V scores would be equal to their true score ( ), the score obtained with no error), and the standard deviation of the distribution of their raw scores would be the conditional standard error. Since there is a one-to-one correspondence between the raw score and in the Rasch mod2 792 QS we can apply this concept more g2 792 n2 792 rall to all students who obtained a particular raw score and calculate the probability of obtaining each possible raw score, JLYHQ WKH VWXGHQW¶V WUXH VFRUH The standard deviation of this conditional distributi

Conditional Standard Error of Measurement Characteristics

The relationship between the scale score CSEM and depends both on the nature of the raw-to-scale score transformation (Kolen and Brennan, 2005; Kolen and Lee, 2011) and on whether the CSEM is derived from the raw scores or from (Lord, 1980). The pattern of CSEMs for raw scores and linear transformations of the raw score tend to have a characteristic ³ L Q Y H U W H G D S H Z L W K V P D O O H U D W W K H H Q G of the score continuum and larger CSEMs toward the middle of the distribution.

Achievable raw score points for these distributions are spaced equally across the score U D Q J H . R O H Q D Q G % U H Q Q D Q S V W D W H ³ : K H Q



Several factors might affect decision consistency and accuracy. One important factor is the reliability of the scores. All other things being equal, more reliable test scores tend to result in more similar reclassifications and less measurement error. Another factor is the location of the cut score in the score distribution. More consistent and accurate classifications are observed when the cut scores are located away from the mass of the score distribution. The number of performance levels is also a consideration. Consistency and accuracy indices based on four performance levels should be lower than those based on two performance levels. This is not surprising, since classification and accuracy using four performance levels would allow more opportunity to change performance levels. Hence, there would be more classification errors and less accuracy with four performance levels, resulting in lower consistency indices.

Results and Observations The results for the dichotomies created by the three cut scores, are presented in Table 8. The tabled values are derived with the program *BB-Class* (Brennan, 2004) using the Livingston and Lewis method. Decision consistency ranged from 0.89 to 0.94, and the decision accuracy ranged from 0.92 to 0.96. Both decision consistency and accuracy values based on individual cut points indicate very good consistency and accuracy of examinee classifications, as shown in Table 8.

Table 8 Decision Consistency and Accuracy Results: Regents Examination in Physical Setting/Earth Science

Statistic	1/2	2/3	3/4
Consistency	0.94	0.91	0.89
Accuracy	0.96	0.94	0.92

4.4

Tabl Group Means: Regents Examination in Physical Setting/Earth Science

The Regents Examination in Physical Setting/Earth Science measures student achievement on the NYS P ±12 Curriculum Standards for Physical Setting/Earth Science. The Physical Setting/Earth Science standards can be found at: http://www.nysed.gov/curriculum-instruction/science/.

Content Validity

Content validity is necessarily concerned with the proper definition of the construc



Figure E.1 Scatter Plot: Regents Examination in Physical Setting/Earth Science

Table E.3 Descriptive Statistics in p-value and Point-Biserial Correlation: Regents Examination in Physical Setting/Earth Science

Statistics	N	Mean	Min	Q1	Median	Q3	Max
p-value	86	0.41	0.14	0.30	0.40	0.53	0.80
Point-Biserial	86	0.31	0.13	0.23	0.31	0.38	0.67

Figure E.2 Student Performance Map: Regents Examination in Physical Setting/Earth Science



Figure E.3 Scree Plots: Regents Examination in Physical Setting/Earth Science

Table E.3 Summary of Item Residual Correlations: Regents Examination in Physical Setting/Earth Science

Statistic Type	Value
N	3570
Mean	> .01
SD	0.03
Minimum	> .10
P_{10}	> .05
P_{25}	> .03
P_{50}	> .01
P_{75}	0.00
P_{90}	0.02
Maximum	0.26
>|0.20|	2

Table E.8 Group Means: Regents Examination in Physical Setting/Earth Science